

# On Geodesic Flow and Energy Functional on Riemannian Manifolds

Yang Liu

Shenzhen Technology University, Shenzhen, China

**Email address:**

yliu1mat@gmail.com

**To cite this article:**

Yang Liu. On Geodesic Flow and Energy Functional on Riemannian Manifolds. *Pure and Applied Mathematics Journal*.

Vol. 10, No. 5, 2021, pp. 104-106. doi: 10.11648/j.pamj.20211005.11

**Received:** August 20, 2021; **Accepted:** September 2, 2021; **Published:** October 12, 2021

**Abstract:** In this paper, we study the geodesic flow and the energy functional on a Riemannian manifold and show that the geodesics have minimal energy, in other words, are the minimizers of the energy functional, from the new perspective of involution, and that the geodesic flow is a Hamiltonian flow which has a close connection with the canonical symplectic structure on the tangent bundle of a Riemannian manifold.

**Keywords:** Geodesic Flow, Energy Functional, Hamiltonian Flow, Vector Bundle

## 1. Introduction

It is well known that geodesics are the shortest lines connecting points in manifolds equipped with metrics, on which one can see, for instance, the researches [3, 13, 2, 16, 7], and on the other hand, geodesics can also be viewed as minimal hypersurfaces of two-dimensional Riemannian manifolds. Further, in the next section, we will show that geodesic flows on a Riemannian manifold are Hamiltonian flows of the energy functional, and it has a close connection with the canonical symplectic structure on the tangent bundle of Riemannian manifolds. On the other hand, the related complex structure in the differential aspect was studied in many researches [17, 11, 10], [12, 14, 4]. Recently, it has been found that the symplectic measurements induced by a symplectic form can potentially be used to address problems in convex geometry, on which one can see, for example, the researches [1, 15, 6, 8, 5]. In this paper, we study the geodesic flow and the energy functional on a Riemannian manifold. The main contribution of this paper is to show that the geodesics have minimal energy, in other words, are minimizers of the energy functional, from the novel perspective of involution, and that the geodesic flow is a Hamiltonian flow, which has a close connection with the canonical symplectic structure on the tangent bundle of a Riemannian manifold.

## 2. Geodesic Flow and Energy Functional

Let  $M$  be an  $n$ -dimensional Riemannian manifold with metric  $g$ ,  $\xi_x$  be any vector in the tangent bundle  $TM$ , and  $c(t)$  be a curve in  $TM$  starting at  $c(0) = \xi_x$  with initial velocity

$$c'(0) = (X, \Xi) \in T_{\xi_x} TM, \quad (1)$$

then

$$(X, \Xi) \in H_{\xi_x} TM, \quad (2)$$

Where  $H_{\xi_x} TM$  is the horizontal space, for which one can see Shoshichi Kobayashi and Katsumi Nomizu's study [9], if and only if  $\nabla_{\xi_x} X = 0$ , where  $H_{\xi_x} TM$  is the horizontal space of the tangent bundle  $TM$ .

In terms of coordinates  $(x^1, \dots, x^n, \xi^1, \dots, \xi^n)$ , let

$$X := \sum_{i=1}^n X_i \frac{\partial}{\partial x^i} \quad (3)$$

and

$$\Xi := \sum_{i=1}^n \Xi_i \frac{\partial}{\partial \xi^i}. \quad (4)$$

Then we have

$$\Xi_k + \sum_{i,j=1}^n \Gamma_{ij}^k X_i \xi_j = 0 \quad (5)$$

by  $\nabla_{\xi_x} X = 0$ .

On the other hand, for vertical space, we have that  $(X, \Xi) \in V_{\xi_x} TM$  holds if and only if  $X = 0$ , i.e.  $X_i = 0$  for all  $i$ .

Here is a direct way to show that a geodesic flow is a Hamiltonian flow of the energy functional. In terms of

coordinates, we know that for the geodesic vector field

$$\chi(\xi_x) := \sum_{k=1}^n \xi_k \frac{\partial}{\partial x^k} - \sum_{k=1}^n \sum_{i,j=1}^n \Gamma_{ij}^k \xi_i \xi_j \frac{\partial}{\partial \xi^k}, \quad (6)$$

on  $TM$  by [5], we have

$$\begin{aligned} i_{\chi(\xi_x)} d\alpha_{\xi_x}(\cdot) &= d\alpha_{\xi_x}(\chi(\xi_x), \cdot) \\ &= dg_{(\xi_x)} \left( \xi_x, \pi_* \left( \sum_{k=1}^n \xi_k \frac{\partial}{\partial x^k} - \sum_{k=1}^n \sum_{i,j=1}^n \Gamma_{ij}^k \xi_i \xi_j \frac{\partial}{\partial \xi^k} \right) \right) (\cdot) \\ &= dg(\xi_x, \xi_x)(\cdot) \\ &= dE_{\xi_x}(\cdot) \end{aligned} \quad (7)$$

So if  $\omega := d\alpha$  is the natural symplectic form on  $TM$ , we then have the following lemma.

Lemma 1. Geodesic flows on a Riemannian manifold are Hamiltonian flows of the energy functional. More precisely,  $i\chi\omega = dE$ .

There are some relations between the energy functional and the canonical differential forms on Riemannian manifolds. We first consider the case  $\Gamma_{ij}^k = 0$ , and in this case it means  $\Xi_k = 0$  for all  $k$  by [5] if  $(X, \Xi) \in H_{\xi_x} TM$ . There is a natural involution between the horizontal space and the vertical space, which is

$$\begin{aligned} \iota : H_{\xi_x} TM &\rightarrow V_{\xi_x} TM \\ \iota \left( \sum k_i \frac{\partial}{\partial x^i} \right) &= \sum k_i \frac{\partial}{\partial \xi^i}. \end{aligned} \quad (8)$$

Thus

$$\begin{aligned} \alpha_{\xi_x} \circ \iota \left( \sum_{i=1}^n \left( k_i \frac{\partial}{\partial \xi^i} \right) \right) &= \alpha_{\xi_x} \left( \sum_{i=1}^n k_i \frac{\partial}{\partial x^i} \right) \\ &= g \left( \xi_x, \pi_* \left( \sum_{i=1}^n \left( k_i \frac{\partial}{\partial x^i} \right) \right) \right) \\ &= \sum_{i,j=1}^n g_{ij} \xi_i \xi_j \end{aligned} \quad (9)$$

$$\begin{aligned} \alpha_{\xi_x} \circ \iota \left( \sum_{i=1}^n \left( k_i \frac{\partial}{\partial \xi^i} \right) \right) &= \alpha_{\xi_x} \left( \sum_{i=1}^n k_i \frac{\partial}{\partial x^i} - \sum_{k=1}^n \sum_{i,j=1}^n \Gamma_{ij}^k k_i \xi_j \frac{\partial}{\partial \xi^k} \right) \\ &= g \left( \xi_x, \pi_* \left( \sum_{k=1}^n k_i \frac{\partial}{\partial x^i} - \sum_{k=1}^n \sum_{i,j=1}^n \Gamma_{ij}^k k_i \xi_j \frac{\partial}{\partial \xi^k} \right) \right) \\ &= \sum_{i,j=1}^n g_{ij} \xi_j k_j \end{aligned} \quad (13)$$

and

$$\begin{aligned} dE_{\xi_x} \left( \sum_{i=1}^n k_i \frac{\partial}{\partial \xi^i} \right) &= \sum_{i=1}^n k_i \frac{\partial}{\partial \xi^i} E(\xi) \\ &= \sum_{i=1}^n k_i \frac{\partial}{\partial \xi^i} g(\xi, \xi) \\ &= \sum_{i,j=1}^n g_{ij} \xi_j k_j. \end{aligned} \quad (14)$$

Hence, it follows that

$$\alpha_{\xi_x} \circ \iota \left( \sum_{i=1}^n \left( k_i \frac{\partial}{\partial \xi^i} \right) \right) = dE_{\xi_x} \left( \sum_{i=1}^n k_i \frac{\partial}{\partial \xi^i} \right). \quad (15)$$

In other words, we have proved the following theorem:

Theorem 2. Let  $E$  be the energy functional on the Riemannian manifold  $M$  and let  $\alpha$  be the natural 1-form on  $TM$  whose differential is the natural symplectic form on  $TM$ . Then we have

On the other hand, for  $dE$ , we have

$$\begin{aligned} dE_{\xi_x} \left( \sum_{i=1}^n k_i \frac{\partial}{\partial \xi^i} \right) &= \sum_{i=1}^n k_i \frac{\partial}{\partial \xi^i} E(\xi_x) \\ &= \sum_{i=1}^n k_i \frac{\partial}{\partial \xi^i} g(\xi_x, \xi_x) \\ &= \sum_{i,j=1}^n g_{ij} \xi_i k_j \end{aligned} \quad (10)$$

Therefore

$$dE_{\xi_x}((0, \Xi)) = \alpha_{\xi_x} \circ \iota((0, \Xi)) \quad (11)$$

for any  $(0, \Xi) \in V_{\xi_x} TM$ .

Let us now consider the general case. For any  $(X, \Xi) \in H_{\xi_x} TM$ , we have  $\Xi_k = -\sum_{i,j=1}^n \Gamma_{ij}^k \chi_i \xi_j$ , analogous to the case  $\Gamma_{ij}^k = 0$ , an involution we can have is

$$\begin{aligned} \iota : V_{\xi_x} TM &\rightarrow H_{\xi_x} TM \\ \iota \left( \sum k_i \frac{\partial}{\partial \xi^i} \right) &= \sum k_i \frac{\partial}{\partial x^i} - \sum \Gamma_{ij}^k k_i \xi_j \frac{\partial}{\partial \xi^k}. \end{aligned} \quad (12)$$

Therefore,

$$dE = \alpha \circ \iota \quad (16)$$

on the vertical bundle of  $TTM$ .

Remark 3. Since both sides on the horizontal bundle of  $TTM$  are zeros, then  $dE = \alpha \circ \iota$  on  $TTM$ .

### 3. Conclusion

The geodesics have minimal energy, in other words, are

the minimizers of the energy functional, from the new perspective of involution, and that the geodesic flow is a Hamiltonian flow which has a close connection with the canonical symplectic structure on the tangent bundle of a Riemannian manifold.

## Declaration of Interest

The author declares that there is no conflict of interest.

## Data Availability Statement

The author confirms that the data supporting the findings of this study are available within the article or its supplementary materials.

## Acknowledgements

This work is supported in part by a research grant from Shenzhen Municipal Finance.

## References

- [1] Naeem Alkoumi and Felix Schlenk. Shortest closed billiard orbits on convex tables. *Manuscripta Mathematica*, 147 (3): 365–380, 2015.
- [2] Georgios Arvanitidis, Soren Hauberg, Philipp Hennig, and Michael Schober. Fast and robust shortest paths on manifolds learned from data. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1506–1515. PMLR, 2019.
- [3] Luther Pfahler Eisenhart. *Riemannian geometry*. Princeton university press, 1997.
- [4] Wei Guo Foo and Joel Merker. Differential e-structures for equivalences of 2-nondegenerate levi rank 1 hypersurfaces m5 in c3. *arXiv preprint arXiv: 1901.02028*, 2019.
- [5] Guilherme França, Alessandro Barp, Mark Girolami, and Michael I Jordan. Optimization on manifolds: A symplectic approach. *arXiv preprint arXiv: 2107.11231*, 2021.
- [6] Hajime Fujita, Yu Kitabepu, and Ayato Mitsuishi. Distance functions on convex bodies and symplectic toric manifolds. *arXiv preprint arXiv: 2003.02293*, 2020.
- [7] Jürgen Jost and Jèurgen Jost. *Riemannian geometry and geometric analysis*, volume 42005. Springer, 2008.
- [8] Roman Karasev. Mahlers conjecture for some hyperplane sections. *Israel Journal of Mathematics*, 241 (2): 795–815, 2021.
- [9] Shoshichi Kobayashi and Katsumi Nomizu. *Foundations of differential geometry*, volume 1. New York, 1963.
- [10] Yang Liu. On the range of cosine transform of distributions for torusinvariant complex minkowski spaces. *Far East Journal of Mathematical Sciences (FJMS)*, 39 (2): 137–157, 2010.
- [11] Yang Liu. On the lagrangian subspaces of complex minkowski space. *J. Math. Sci. Adv. Appl*, 7 (2): 87–93, 2011.
- [12] Yang Liu. On the kähler form of complex lp space and its lagrangian subspaces. *Journal of Pseudo-Differential Operators and Applications*, 6 (2): 265–277, 2015.
- [13] Peter Petersen, S Axler, and KA Ribet. *Riemannian geometry*, volume 171. Springer, 2006.
- [14] Vandana Rani and Jasleen Kaur. On the structure of lightlike hypersurfaces of an indefinite kaehler statistical manifold. *Differential Geometry-Dynamical Systems*, 23: 229–242, 2021.
- [15] Felix Schlenk. Symplectic embedding problems, old and new. *Bulletin of the American Mathematical Society*, 55 (2): 139–182, 2018.
- [16] Fengzhen Tang, Mengling Fan, and Peter Tiño. Generalized learning riemannian space quantization: A case study on riemannian manifold of spd matrices. *IEEE transactions on neural networks and learning systems*, 32 (1): 281–292, 2020.
- [17] Edoardo Vesentini. Complex geodesics. *Compositio mathematica*, 44 (1-3): 375–394, 1981.